

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



18/01
SGI



PCT/GB 00 / 02361

4



INVESTOR IN PEOPLE

10/031125

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 10 AUG 2000

WIPO PCT

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)


G800/2361

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 6 July 2000


THIS PAGE BLANK (USPTO)

13 JUL 1999

14 JUL 99 E461848-1 D01348
P01/7700 0.00 - 9916434.5

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road

Newport

Gwent NP9 1RH

1. Your reference

G.1344

2. Patent application number

(The Patent Office will fill in this part)

9916434.5

3. Full name, address and postcode of the or of each applicant (underline all surnames)

ELI LILLY AND COMPANY LIMITED,
KINGSCLERE ROAD,
BASINGSTOKE,
HAMPSHIRE RG21 6XA

Patents ADP number (if you know it)

6106002

If the applicant is a corporate body, give the country/state of its incorporation

GREAT BRITIAN

07755 291001

4. Title of the invention

PHARMACEUTICAL COMPOUNDS

5. Name of your agent (if you have one)

C. M. HUDSON

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

LILLY RESEARCH CENTRE,
ERL WOOD MANOR,
WINDLESHAM,
SURREY GU20 6PH

Patents ADP number (if you know it)

2303001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

YES

a) any applicant named in part 3 is not an inventor, or

b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

29 /

Claim(s)

4 /

Abstract

2 /

Drawing(s)

-

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

1 /

Request for preliminary examination and search (Patents Form 9/77)

1 /

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. —

I/We request the grant of a patent on the basis of this application.

Signature

Date

13th July 1999

12. Name and daytime telephone number of person to contact in the United Kingdom

MR. C. M. HUDSON (01276) 853443

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

DUPLICATE

PHARMACEUTICAL COMPOUNDS


This invention relates to novel chemical compounds and their use as pharmaceuticals.

5

It is well known that chemical compounds which modulate the activity of neuronal calcium channels are potentially useful in treating disorders of the central nervous system.

10

The compounds of the invention have the following general formula:



(I)

15

in which the aminosulfonyl group is attached at the 3-
or 4-position, and in which

R¹ is hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀
5 cycloalkyl-C₁₋₄ alkyl or optionally substituted phenyl-
C₁₋₄ alkyl,

R² is C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-
C₁₋₄ alkyl, optionally substituted phenyl-C₁₋₄ alkyl or
10 -(CH₂)₂NR⁵R⁶ where R⁵ and R⁶ are each hydrogen or C₁₋₆
alkyl, and

R³ and R⁴ are each C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀
cycloalkyl-C₁₋₄ alkyl, C₃₋₆ alkenyl, optionally
15 substituted phenyl or optionally substituted phenyl-C₁₋₄
alkyl,

or R¹ and R², or R³ and R⁴, or R⁵ and R⁶, together with
the nitrogen atom to which they are attached, form a
20 carbocyclic group containing 4 to 7 carbon atoms
optionally substituted with one to three methyl or ethyl
groups and optionally containing an oxygen atom or a
further nitrogen atom, said carbocyclic group being

optionally fused to an optionally substituted phenyl group;

or a salt thereof.

5

The compounds of the invention have been found to be active in tests that show modulation of voltage-dependent calcium channels, and are thus indicated for use in the treatment of diseases in which such modulation is beneficial, in particular diseases of the central nervous system.

In the above formula (I), a C₁₋₆ alkyl group includes methyl, ethyl, propyl, isopropyl, butyl, tert. butyl and hexyl, and is preferably methyl or ethyl. A substituted phenyl group is phenyl substituted with one or more, for example one to three, substituents selected from, for example C₁₋₄ alkyl, especially methyl, C₁₋₄ alkoxy, especially methoxy and ethoxy, hydroxy, nitro, cyano, halo, especially chloro or fluoro, trihalomethyl, especially trifluoromethyl, carboxy and C₁₋₄ alkoxy-carbonyl. A halo atom is preferably chlorine, bromine or fluorine. A substituted phenyl group preferably has one to three substituents selected from hydroxy, C₁₋₄ alkyl, halo, nitro and trifluoromethyl. An optionally

substituted phenyl-C₁₋₄ alkyl group is preferably of the formula R-(CH₂)_n- where R is optionally substituted phenyl and n is 1 to 4, but the linking chain can also be branched alkylene. A C₃₋₁₀ cycloalkyl group is
5 preferably, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl and these groups may optionally be substituted by one or two C₁₋₄ alkyl, especially methyl, substituents. A C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl group is one such cycloalkyl group attached
10 to a C₁₋₄ alkyl, and is preferably of the formula R-(CH₂)_n- where R is cycloalkyl and n is 1 to 4. When R³ or R⁴ is C₁₋₆ alkyl it is preferably C₃₋₆ alkyl.

The groups R¹ and R², R³ and R⁴, and R⁵ and R⁶, can form
15 a carbocyclic ring with the nitrogen to which they are attached and optionally also contain an oxygen atom or an additional nitrogen. Preferred examples, including the nitrogen of the amino sulfonyl group, are pyrrolidino, piperazino, morpholino and especially
20 3,5-dimethylpiperidino.

A particular group of compounds of the invention is one of formula (I) in which R¹, R², R³ and R⁴ are each C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl or

optionally substituted phenyl-C₁₋₄ alkyl, and R¹ can in addition be hydrogen, or R¹ and R², or R³ and R⁴ together with the nitrogen atom to which they are attached, form a carbocyclic group as defined above.

5

In a preferred group of compounds R¹, R², R³ and R⁴ are each C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl or optionally substituted phenyl-C₁₋₄ alkyl, and R¹ is in addition hydrogen.

10

It is preferred that R¹ is hydrogen. Furthermore, R³ and R⁴, which can be the same or different, are preferably C₁₋₄ alkyl. It is further preferred that R² is optionally substituted phenyl-C₁₋₄ alkyl.

15

A further preferred group of compounds is one of formula (I) in which R² is $-(CH_2)_2NR^5R^6$.

A further preferred group of compounds is one of
20 formula (I) in which R³ or R⁴ is C₃₋₆ alkyl or when R³ and R⁴ are taken together with the nitrogen atom they form a piperidine ring which is substituted at the 3- and/or 5-positions with one or two methyl or ethyl substituents.


It will be appreciated that the compounds of the invention can contain one or more asymmetric carbon atom which gives rise to enantiomers. The compounds can be
5 prepared as racemates or can be made from enantiomeric intermediates. Both racemates and enantiomers form part of the present invention.

It will also be understood that salts of the compounds
10 of the invention can be prepared and such salts are included in the invention. They can be any of the well known acid addition salts. Acid addition salts are preferably the pharmaceutically acceptable non-toxic addition salts with suitable acids, such as those with
15 inorganic acids, for example hydrochloric, hydrobromic, nitric, sulfuric or phosphoric acids, or with organic acids, such as organic carboxylic acids, for example glycollic, maleic, fumaric, malic, oxalic, tartaric, citric, salicylic or o-acetoxybenzoic acids, or organic
20 sulfonic acids, methane sulfonic, 2-hydroxyethane sulfonic, toluene-p-sulfonic or naphthalene-2-sulfonic acids.

In addition to pharmaceutically-acceptable salts, other
25 salts are included in the invention. They may serve as intermediates in the purification of compounds or in the

preparation of other, for example pharmaceutically-acceptable, salts, or are useful for identification, characterisation or purification.

- 5 The invention includes a process for producing the compounds of formula (I) above which comprises reducing a compound of the fomula:




(II)

10

The reaction is preferably carried out in an organic solvent, for example, at a temperature of 0° C. to 100° C., employing a reducing agent, for example lithium aluminium hydride.

15

Compounds of formula (II) can readily be prepared by conventional methods, for example, by reacting a compound of the formula:



(III)


5

where X is a leaving group such as, for example, halo or hydroxy, with an amine of the formula HNR¹R².

- 10 The reaction is preferably carried out in an organic solvent such as, for example, chloroform or acetonitrile, at a temperature of from 0° C. to 100° C. such as, for example, ambient temperature.
- 15 Intermediate compounds of formula (III) are known in the art and can be readily prepared by known methods. When an acid halide is employed (X is halo such as, for example, chloro), the reaction is preferably carried out

in the presence of a solid phase scavenger to absorb the acid liberated by the reaction. When the free acid is employed (X is hydroxy), a condensing reagent such as, for example, dimethylaminopropyl-ethylcarbodiimide can
5 be employed.

A further route to the compounds of the invention, which is also included in the invention, involves the reduction of the imine corresponding to the compound of
10 formula (III):



employing a reducing agent as, for example, sodium
15 borohydride. Compounds of formula (IV) can readily be prepared by reacting an amine of formula R¹R²NH with the

appropriate benzaldehyde derivative, which can, in its turn, be prepared by reducing the corresponding benzoic acid derivative to the alcohol, followed by oxidation to the required benzaldehyde intermediate.

5

Amine reactants of the formula HNR^1R^2 are well known and can be readily prepared by known methods. Those in which R^2 is $-(\text{CH}_2)_2\text{NR}^5\text{R}^6$ can, for example, be prepared by reductive amination, that is, by reacting the
10 appropriate diamine with an aldehyde in reducing conditions.

Alternatively, compounds of formula (I) in which R^2 is $-(\text{CH}_2)_2\text{NR}^5\text{R}^6$ can be prepared by alkylation of the
15 corresponding compound of formula (I) in which R^1 is hydrogen.

As mentioned above, the compounds of the invention are
20 active in tests that indicate their utility in the treatment of diseases of the central nervous system. The compounds modulate the activity of calcium channels and, in particular, they block voltage sensitive calcium channels as determined in a test based on Boot J. R.,
25 et al., Specificity of autoantibodies in the Lambert-

Eaton Myasthenic Syndrome, Ann NY Acad. Sci. (1997), in which measurements of calcium flux using calcium sensitive dyes are made. Compounds described in the following Examples were found to inhibit voltage-
5 dependent calcium channels in cloned human cell lines expressing specific voltage-dependent calcium channels with an IC₅₀ of less than 10 μ M.

The compounds of the invention are thus indicated for
10 use in the treatment of anoxia, ischaemia, stroke and heart failure, migraine, diabetes, cognitive impairment, pain, epilepsy, traumatic head or spinal injury, AIDS related dementia and blindness, amnesia, neurodegenerative diseases such as Alzheimer's,
15 Parkinson's and Huntington's diseases and age-related memory disorders, Down's syndrome, mood disorders, drug or alcohol addition withdrawal, nausea from chemotherapy, and carbon monoxide or cyanide poisoning.

20 The invention also includes a pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier in association with the compound of the invention or a pharmaceutically acceptable salt or ester thereof.

25

The compound may be administered by various routes, for example by the oral or rectal route, topically or parenterally, for example by injection or infusion, being usually employed in the form of a pharmaceutical composition. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, and/or enclosed within a carrier which may, for example, be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition may be in the form of tablets, lozenges, sachets, cachets, elixirs, suspensions, ointments containing, for example, up to 10% by weight of the compound, soft and hard gelatin capsules, suppositories, injection solutions and suspensions and sterile packaged powders.

Some examples of suitable carriers are lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propyl-hydrobenzoate, talc magnesium stearate and mineral oil.

The compositions of the injection may, as is well known in the art, be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

5

Where the compositions are formulated in unit dosage form, it is preferred that each unit dosage form

contains from 5 mg to 500 mg. The term 'unit dosage form' refers to physically discrete units suitable as

10 unit dosages for human subjects and animals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

15 The active compound is effective over a wide dosage range and, for example, dosages per day will normally fall within the range of from 0.5 to 300 mg/kg, more usually in the range of from 5 to 100 mg/kg. However, it will be understood that the amount administered will
20 be determined by the physician in the light of the relevant circumstances including the conditions to be treated, the choice of compound to be administered and the chosen route of administration, and therefore the above dosage ranges are not intended to limit the scope
25 of the invention in any way.

The invention is illustrated by the following
Preparations and Examples.

5 EXAMPLE 1

4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid

To a stirred solution of di-n-propylamine (3.03 g,
10 0.03 mole) in dry tetrahydrofuran (20 ml) at 0° C.
(ice/salt bath), was added 4-chlorosulfonylbenzoic acid
(2.2 g, 0.01 mole). Stirring was continued for 1 hour.
Ice water was added cautiously and the reaction made
acid with 2NHCl. The 4-[(N,N-di-n-
15 propylamino)sulfonyl]-benzoic acid was collected by
filtration as a white solid which was dried *in vacuo* at
40° C.

20 EXAMPLE 2

4-[(N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-
benzamide

25 To a solution of 4-[(N,N-di-n-propylamino)sulfonyl]-
~~benzoic acid (2.85 g, 0.01 mole) in dry dichloromethane~~

(ml) at 0° C. was added oxalyl chloride (2.54 g, 0.02 mole) and dimethylformamide (4 drops). The reaction mixture was stirred for 2 hours. The reaction was evaporated to dryness *in vacuo*. The resulting acid chloride was added to a stirred solution of *p*-methoxybenzylamine (1.51 g, 0.011 mole) and triethylamine (1.11 g, 0.011 mole) in dry tetrahydrofuran (25 ml) at 0-5° C. After stirring for 4 hours the reaction was poured into ice water and extracted with ethyl acetate. The solvent was washed with brine, dried and evaporated to dryness *in vacuo*. Chromatography on flash silica using 10% ethyl acetate/dichloromethane gave 4-[(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide as a white solid. M.p. 132-134° C.

EXAMPLE 3

N,N-di-n-propyl-4-[[(4-methoxybenzyl) amino]methyl] benzenesulfonamide

To a stirred solution of 4-[(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide (1.87g, 4.62mmole) in dry ether (50ml) was added a solution of 2M lithium aluminium hydride in tetrahydrofuran (4.63ml,

9.24mmole). The reaction was heated at reflux for 2 hours. After cooling to room temperature water (1ml) was added dropwise with caution followed by 2N NaOH (1ml). When gas evolution ceased the reaction mixture
5 was filtered through a pad of celite which was well washed with ether. After removal of the solvent *in vacuo* the product was purified by chromatography on flash silica eluting with 10% methanol/ethyl acetate. The resulting amine was converted to the maleic acid
10 salt and re-crystallised from ethanol/ether to give *N,N*-di-*n*-propyl-4-{[(4-methoxybenzyl)amino]methyl} benzenesulfonamide maleate. mp. 133-135°C

Similarly prepared were:

15

N,N-di-*n*-propyl-3-{[(4-methoxybenzyl)amino]methyl} benzenesulfonamide maleate. mp. 160-162°C

N,N-di-*n*-propyl-4-{[(3,4-dimethoxyphenethyl)amino]methyl} benzenesulfonamide maleate. mp. 130-132°C

20 *N*-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)amino]methyl} benzenesulfonamide maleate. mp. 169-171°C

N-(3,3-dimethylpiperidino)-4-{[(4-fluorobenzyl)amino]methyl} benzenesulfonamide maleate. mp. 196-198°C

25 *N,N*-di-*n*-propyl-3-{[(4-fluorobenzyl)amino]methyl} benzenesulfonamide maleate. mp. 168-170°C

N-phenyl-*N*-*n*-propyl-4-{[(dimethylamino)amino]methyl}

benzenesulfonamide maleate. mp. 154-156°C

N-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)-*N*-methylamino]methyl}benzenesulfonamide maleate. Mass spectrum: $MH^+ = 405$ (TSP+)

5 *N*-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)-*N*-benzylamino]methyl}benzenesulfonamide maleate. mp. 183-185°C

N-phenyl-*N*-methyl-3-{[(4-fluorobenzyl)amino]methyl}benzenesulfonamide maleate. mp. 194-196°C

10 *N*-phenyl-*N*-*n*-butyl-4-{[hexylamino]methyl}benzenesulfonamide maleate. mp. 106-108°C

N-(3-ethylpiperidino)-3-{[(4-fluorobenzyl)amino]methyl}benzenesulfonamide maleate. mp. 140-142°C

N-(3,3-dimethylpiperidino)-3-{[(cyclohexylmethyl)amino]methyl}benzenesulfonamide hydrochloride. mp. 147-149°C

N-(3-methylpiperidino)-4-{[(4-chlorophenethyl)amino]methyl}benzenesulfonamide maleate. mp. 176-178°C

N-(3-methylpiperidino)-4-{[(4-chlorophenethyl)-*N*-methylamino]methyl}benzenesulfonamide maleate. mp. 168-170°C

3-{[[2-(dimethylamino)ethyl](4-fluorobenzyl)amino]methyl}-*N*-3,3-dimethylpiperidino-benzenesulfonamide maleate as an oil. Mass

25 spectrum($MH^+ = 462$ (10%)) (TSP+)

3-{[[2-(dimethylamino)ethyl](cyclohexylmethyl)
amino]methyl}-N-3,3-dimethylpiperidino-
benzenesulfonamide maleate. mp. 149-151°C

5

EXAMPLE 4

4-{[[2-(piperidino)ethyl](2-[3,4-
dimethoxy]phenylethyl)amino]methyl}-N,N-di-n-
10 propylbenzene sulfonamide dihydrochloride

To solution of N,N-di-n-propyl-4-{[(3,4-
dimethoxyphenethyl)amino]methyl}benzene sulfonamide (550
mg, 1mmole) in dry acetonitrile (100ml) was added sodium
15 carbonate (440mg, 4.4mmole), potassium iodide (166mg,
1mmole) and 2-chloroethylpiperidine hydrochloride
(184mg, 1mmole). The reaction was stirred and heated at
reflux for 18 hours. The reaction was poured into ice
water and extracted with ethyl acetate, washed with
20 brine, dried and evaporated to dryness *in-vacuo*.
Chromatography on flash silica by elution with
10%methanol/dichloromethane gave 4-{[[2-
(piperidino)ethyl](2-[3,4-dimethoxy]phenylethyl)amino]
methyl}-N,N-di-n-propylbenzenesulfonamide which was
25 crystallised as its dihydrochloride salt. mp. 135-137°C

EXAMPLE 5

N-(3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl)-*N*-(4-methylbenzyl)amine

A mixture of a 0.15 M solution of 3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in methanol (0.25 ml) and a 0.1 M solution of 4-methylbenzylamine in methanol (0.25 ml) was stirred at room temperature for 1 hour. A 0.15 M solution of sodium borohydride in methanol (0.25 ml) was added and stirring continued for a further 16 hours. The mixture was then applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a 2 M solution of ammonia in methanol (2.5 ml). The eluate was vacuum evaporated to give the required product. (TS-MS: m/z 387, [M+H]⁺).

The following compounds were similarly prepared (mass spectrum values are given in brackets).

N-(3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl)-*N*-(3-(4-methylpiperazin-1-yl)propyl)amine (423)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-(3-morpholin-4-ylpropyl)amine (410)

5 *N*-(4-chlorobenzyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (407/408)

N-(cyclohexylmethyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (379)

10 *N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-[3-(1*H*-imidazol-1-yl)propyl]amine (391)

N-butyl-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (339)

15

N-(*tert*-butyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (339)

20 *N*-(2-chlorobenzyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (407/408)

N-(4-chlorophenethyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (421/422)

N-(2-chlorophenethyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (421/422)

5 *N*-(2,4-dichlorobenzyl)-*N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (442)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-isopentylamine (353)

10 *N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-(3-methoxypropyl)amine (355)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-(2-methylbenzyl)amine (387)

15

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-(3-methylcyclohexyl)amine (379)

20 *N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-hexylamine (367)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-propylamine (325)

25 *N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-(4-methylphenethyl)amine (401)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-[3-(trifluoromethyl)benzyl]amine (441)

5 *N*-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-*N*-[3-(trifluoromethyl)phenethyl]amine (455)

EXAMPLE 6

10 1-({3-[(4-benzylpiperidin-1-yl)methyl]phenyl}sulfonyl)-3,3-dimethylpiperidine

A mixture of a 0.15 M solution of 3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in
15 dichloromethane (0.25 ml), a 0.1 M solution of 4-benzylpiperidine in dichloromethane (0.25 ml) and a 0.15 M solution of sodium tri-acetoxyborohydride in dichloromethane (0.25 ml) was stirred at room temperature for 22 hours. Methanol (1 ml) was added and
20 the mixture applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a 2 M solution of ammonia in methanol (2.5 ml). The eluate was vacuum evaporated to give the
25 required product. (TS-MS: m/z 441, [M+H]⁺).

The following compounds were similarly prepared (mass spectrum values are given in brackets).

2-(butyl{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amino)ethan-1-ol (383)

2-(benzyl{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amino)ethan-1-ol (417)

10 N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-N,N-bis(2-methoxyethyl)amine (399)

1-(3,4-dichlorophenyl)-4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperazine (497)

15

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-N-ethyl-N-(pyridin-4-ylmethyl)amine (402)

1-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-4-(4-fluorophenyl)piperazine (446)

4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperazine-1-carbaldehyde (380)

25 4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}morpholine (353)

1-[4-(4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperazin-1-yl)phenyl]ethan-1-one
(470)

5

3,3-dimethyl-1-[[3-(pyrrolidin-1-ylmethyl)phenyl]sulfonyl]piperidine (337)

2-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-
10 1,2,3,4-tetrahydroisoquinoline (399)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-N,N-
dipropylamine (367)

15 1-benzhydryl-4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperazine (518)

N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-N-(2-
methoxyethyl)-N-propylamine (383)

20

EXAMPLE 7

1-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperidine-4-carboxamide

25

A mixture of a 0.15 M solution of 3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in methanol (0.25 ml), a 0.1 M solution of piperidine-4-carboxamide in methanol/acetic acid 4:1 v/v (0.25 ml) and a 0.15 M
5 solution of sodium cyanoborohydride in methanol (0.25 ml) was stirred at room temperature for 18 hours. The mixture applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a
10 2 M solution of ammonia in methanol (2.5 ml) and the eluate vacuum evaporated. The residue was dissolved in chloroform (2 ml) and the solution added to isocyanatomethyl-polystyrene (loading 1 mmole/g, 100 mg). The suspension was shaken at room temperature for
15 16 hours, then filtered. The resin was washed with chloroform (2 x 2 ml) and the combined filtrates vacuum evaporated to give the required product. (TS-MS: m/z 394, [M+H]⁺).

20

The following Examples illustrate typical formulations containing a compound of the invention.

25 EXAMPLE 8

Tablets each containing 10 mg of active ingredient are made up as follows:

	Active ingredient	10 mg
5	Starch	160 mg
	Microcrystalline cellulose	100 mg
	Polyvinylpyrrolidone (as 10% solution in water)	13 mg
	Sodium carboxymethyl starch	14 mg
	Magnesium stearate	3 mg
10		_____
	Total	300 mg

The active ingredient, starch and cellulose are mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders and passed through a sieve. The granules so produced are dried and re-passed through a sieve. The sodium carboxymethyl starch and magnesium stearate are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 300 mg..

EXAMPLE 9

Capsules each containing 20 mg of active ingredient are made as follows:

	Active ingredient	20 mg
	Dried starch	178 mg
	Magnesium stearate	2 mg
5		_____
	Total	200 mg

The active ingredient, starch and magnesium stearate are
10 passed through a sieve and filled into hard gelatine
capsules in 200 mg quantities.

EXAMPLE 10

15

Capsules each containing 20 mg of medicament are made as
follows:

	Active ingredient	20 mg
20	Lactose	171 mg
	Sodium lauryl sulphate	2 mg
	Sodium starch glycollate	6 mg
	Magnesium stearate	1 mg

25		200 mg

The active ingredient, lactose, sodium lauryl sulphate and sodium starch glycollate are mixed thoroughly. The blend is mixed with the magnesium stearate and filled
5 into hard gelatine capsules in 200 mg quantities.

EXAMPLE 11

Tablets each containing 20 mg and medicaments are made
10 as follows:

	Active ingredient	20 mg
	Lactose	103 mg
	Microcrystalline cellulose	150 mg
15	Hydroxypropylmethylcellulose	15 mg
	Sodium starch glycollate	9 mg
	Magnesium stearate	3 mg


		300 mg
20		_____

The active ingredient, lactose, microcrystalline cellulose, sodium starch glycollate and hydroxypropylmethylcellulose are passed through a sieve
25 and blended together. Water is added to the blended
~~powders to form a damp mass. The damp mass is passed~~

through a coarse screen, dried, then re-screened. The dried granules are mixed with the magnesium stearate and compressed into tablets of 300 mg weight.

CLAIMS

1. A compound of the formula



(I)

in which the aminosulfonyl group is attached at the
3- or 4-position, and in which

10 R¹ is hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀
cycloalkyl-C₁₋₄ alkyl or optionally substituted
phenyl-C₁₋₄ alkyl,

R² is C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀
15 cycloalkyl-C₁₋₄ alkyl, optionally substituted

phenyl-C₁₋₄ alkyl or -(CH₂)₂NR⁵R⁶ where R⁵ and R⁶ are each hydrogen or C₁₋₆ alkyl, and

5 R³ and R⁴ are each C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, C₃₋₆ alkenyl, optionally substituted phenyl or optionally substituted phenyl-C₁₋₄ alkyl,

10 or R¹ and R², or R³ and R⁴, or R⁵ and R⁶, together with the nitrogen atom to which they are attached, form a carbocyclic group containing 4 to 7 carbon atoms optionally substituted with one to three methyl or ethyl groups and optionally containing an oxygen atom or a further nitrogen atom, said
15 carbocyclic group being optionally fused to an optionally substituted phenyl group;

or a salt thereof.

20 2. A compound according to Claim 1 in which R¹, R², R³ and R⁴ are each C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl or optionally substituted phenyl-C₁₋₄ alkyl, and R¹ can in addition be
~~hydrogen, or R¹ and R², or R³ and R⁴ together with~~

the nitrogen atom to which they are attached, form a carbocyclic group.

3. A compound according to Claim 2 in which R^1 , R^2 , R^3
5 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl, and R^1 can in addition be hydrogen.
- 10 4. A compound according to Claim 3 in which R^1 is hydrogen, R^2 is optionally substituted phenyl- C_{1-4} alkyl and R^3 and R^4 are C_{1-6} alkyl.
- 15 5. A compound according to Claim 1 in which R^2 is $-(CH_2)_2NR^5R^6$.
6. A compound according to Claim 1 or 5 in which R^3 or R^4 is C_{3-6} alkyl or when R^3 and R^4 are taken
together with the nitrogen atom they form a
20 piperidine ring which is substituted at the 3- and/or 5-positions with one or two methyl or ethyl substituents.

7. A pharmaceutical formulation comprising a compound according to any of Claims 1 to 6 or a pharmaceutically acceptable salt thereof, together with a diluent or carrier therefor.

5


8. A compound according to any of Claims 1 to 6, for use as a pharmaceutical.

10

9. Use of a compound according to any of Claims 1 to 6, in the manufacture of a medicament for treating a disease of the central nervous system.

ABSTRACT

A pharmaceutical compound of the formula



5

in which the aminosulfonyl group is attached at the 3- or 4-position, and in which

10 R¹ is hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl or optionally substituted phenyl-C₁₋₄ alkyl,

R² is C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl, optionally substituted phenyl-C₁₋₄ alkyl or

$-(CH_2)_2NR^5R^6$ where R^5 and R^6 are each hydrogen or C_{1-6} alkyl, and

R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10}
5 cycloalkyl- C_{1-4} alkyl, C_{3-6} alkenyl, optionally
substituted phenyl or optionally substituted phenyl- C_{1-4}
alkyl,

or R^1 and R^2 , or R^3 and R^4 , or R^5 and R^6 , together with
10 the nitrogen atom to which they are attached, form a
carbocyclic group containing 4 to 7 carbon atoms
optionally substituted with one to three methyl or ethyl
groups and optionally containing an oxygen atom or a
further nitrogen atom, said carbocyclic group being
15 optionally fused to an optionally substituted phenyl
group;

or a salt thereof.

20

LET NO : G800 / 2361

Form 23/77 : 15.6.00

Agent : El. Lilly & Co Ltd

THIS PAGE BLANK (10/10)
